

# Physical chemistry and radiochemistry

Prof. Krisztina László (463-)18-93

klaszlo@mail.bme.hu

Building: F, Staircase: I, 1st floor, Room 135

<http://oktatas.ch.bme.hu/oktatas/konyvek/fizkem/PHCR>

# Requirements

Weekly contact hrs: 2+0+1

Individual work with regular consultation

2 tests (min. 50 % each) after 6 and 12 weeks  
(exact time to be negotiated)

Next contact: 29 September, 8:15

Building H, ground floor (ring the bell beside the big glass door)

# RADIOCHEMISTRY

- ✓ to understand the nuclear forces acting in the nucleus of the atoms
- ✓ the kinds and source of nuclear radiations
- ✓ interactions of nuclear radiation with the matter
- ✓ applications




Antoine Henri *Becquerel*  
(1852 - 1908)



Maria *Skłodowska-Curie*  
(1867 – 1934)

# The nucleus



after <http://astronomyonline.org/Science/Images/Mathematics/AtomicStructureSmall.jpg>

$$\Delta E = mc^2$$

$$A = Z + N$$

A: mass number


Z: atomic number

	m	E, MeV
p	$1.6726 \times 10^{-24}$ g	938.27
n	$1.6749 \times 10^{-24}$ g	939.55
$e^-$	$9.109 \times 10^{-28}$ g	0.51

# Stable nuclides




$$A = Z + N$$



# Binding energy of the nucleus

$$\Delta E = \Delta mc^2$$

$$M < Zm_p + Nm_n$$



# Classification of the nuclides

Isotope: identical  $Z$

Isobar: identical  $A$

Isotone: identical  $N$

Isotope effect

i Radioactive isotope !

applications

spectroscopies (resonance, MS)

solvent (NMR, neutron scattering)

enrichment of isotopes

CSIA: compound specific isotope analysis

Negligible?

labelling

unortodox organic synthesis routes



# Radioactivity


Spontaneous transformation of the unstable nucleus.

The properties of the nucleus change in time and energy is lost.


All the conservation laws are met.

# Types of radioactive decay

# Isomeric transition



$$\Delta E = h \cdot \nu$$



line spectrum


## Examples

nuclide	$T_{1/2}$	$E_{\gamma}$ , MeV
${}^{60m}\text{Co}$	10.5 min	0.059
${}^{99m}\text{Tc}$	6.0 h	0.143


Z	Nuclide	$T_{1/2}$	Way of decay	Particle energy, MeV		Gamma energy, MeV		$\eta$	Production	$\sigma'$	Daughter
27						2,02	11 %				
						2,60	16 %				
						2,99	1 %				
						3,25	12 %				
						3,47	1 %				
	$^{57}\text{Co}$	270 d	<i>E.X.</i>		100 %	0,014	6 %	83 %	$^{56}\text{Fe}(d,n)$ $^{60}\text{Ni}(p,\alpha)$	0,9	
						0,122	88 %	1 %			
						0,136	10 %	1 %			
	$^{58}\text{Co}$	71,3 d	<i>E.X.</i> $\beta^+$	0,47	85 % 15 %	0,81	100 %		$^{58}\text{Ni}(n,p)$		
						1,62	0,5 %				
						0,51 ( $\beta^+$ )					
	$^{60m}\text{Co}$	10,5 min	<i>I</i>		100 %	0,059	0 %	$\approx 100\%$	$^{59}\text{Co}(n,\gamma)$	19	$^{60}\text{Co}$
	$^{60}\text{Co}$	5,27 a	$\beta^-$	0,31	$\approx 100\%$	1,17	100 %		$^{59}\text{Co}(n,\gamma)$	37	
				1,48	0,01 %	1,33	100 %				
28	$^{63}\text{Ni}$	92 a	$\beta^-$	0,067	100 %				$^{62}\text{Ni}(n,\gamma)$	0,77	
	$^{65}\text{Ni}$	2,521 h	$\beta^-$	0,60	$\approx 23\%$	0,37	5 %		$^{64}\text{Ni}(n,\gamma)$	0,016	
				1,01	$\approx 8\%$	1,11	13 %				
				2,10	$\approx 69\%$	1,49	18 %				
29	$^{64}\text{Cu}$	12,9 h	$\beta^-$ $\beta^+$	0,57	38 %	0,51 ( $\beta^+$ )			$^{63}\text{Cu}(n,\gamma)$	3,0	
				0,66	19 %	1,34	0,6 %				
				<i>E.X.</i>		43 %					
	$^{66}\text{Cu}$	5,10 min	$\beta^-$	0,76	$< 0,2\%$	0,83	0,2 %		$^{65}\text{Cu}(n,\gamma)$	0,56	
				1,59	$\approx 9\%$	1,04	9 %				
				2,63	$\approx 91\%$						

# $\beta$ - decays



$\beta^-$ -decay



$\beta^+$ -decay



Electron capture



common:

$A = \text{constant}$

$\Delta Z = \pm 1$

$\nu$  or  $\tilde{\nu}$

## Examples: pure $\beta^-$ emitters

nuclide	Energia, MeV	$T_{1/2}$
$^3\text{H}$	0.018	12.26 y
$^{14}\text{C}$	0.159	5730 y
$^{32}\text{P}$	1.71	14.3 d
$^{35}\text{S}$	0.167	88 d
$^{90}\text{Sr}$	0.54	28.1 y
$^{90}\text{Y}$	2.25	64 h

## Examples: mixed ( $\beta+\gamma$ ) emitters

nuclide	$T_{1/2}$	$\beta$ -energy, MeV	$\gamma$ -energy, MeV
$^{60}\text{Co}$	5,27 a	0,31	1,17/1,33
$^{131}\text{I}$	8,07 d	0,61	0,36
$^{137}\text{Cs}$	30,23 a	0,51	0,662

## Examples: positron emitters


nuklid	$T_{1/2}$	$E_{\beta^+}$ MeV
$^{11}\text{C}$	20.3 min	0.97
$^{13}\text{N}$	9.97 min	1.2
$^{15}\text{O}$	124 s	1.7
$^{18}\text{F}$	109.7 min	0.064

## Examples: EX (electron capture)


Nuclide	$T_{1/2}$	$E_{\gamma}$ MeV
$^{54}\text{Mn}$	303 d	0.84
$^{125}\text{I}$	60 d	0.035



# $\alpha$ -decay



4-9 MeV



line spectrum

## Example: Alpha emitters

nuclide	$T_{1/2}$
$^{235}\text{U}$	7.1E8 a
$^{226}\text{Ra}$	1600 a
$^{222}\text{Rn}$	3.8 d

## Gamma ray/radiation

Electromagnetic radiation, emitted by the nucleus

Line spectrum

Isomeric transition ("escort" also)

## Beta-radiations

$e^-$  or  $e^+$  radiation coming from the nucleus

Continuous spectrum

May be exclusive (but  $\nu$ !)

May be escorted by gamma or characteristic X-rays

## Alpha-radiation

${}^4_2\text{He}^{2+}$  particles, emitted by the nucleus

Linear spectrum

May be escorted by gamma radiation

# Radioactivity

-Spontaneous decay

-Properties change in time  
chemical identity  
mass

-Energy is released

$h\nu$  from nucleus: gamma-ray  
 $e^-, e^+$  from nucleus: beta-particle  
 ${}^4_2\text{He}^{2+}$  from nucleus: alpha-particle

mass, MeV

typical energy, MeV

-


0.51

~3700

4-9 MeV

Charge!  
spontaneous fission

**Occurs in nature!!!**



# Kinetics of the decay

## Simple decay


$$A \equiv -\frac{dN}{dt} = \lambda N$$

$$N = N_0 e^{-\lambda t} \quad A = A_0 e^{-\lambda t}$$

$$T_{1/2} = \frac{\ln 2}{\lambda} \quad [A] = \frac{1}{\text{time}}$$

$$\frac{1 \text{ decay}}{\text{second}} = 1 \text{ becquerel} = 1 \text{ Bq}$$

$$1 \text{ Ci} = 3.7 \times 10^{10} \text{ Bq}$$



$$I = k\eta A$$

## Radiocarbon dating (or simply carbon dating)

radiometric dating technique based on the decay of  $^{14}\text{C}$  to estimate the age of organic materials (wood, leather, etc.) up to 58,000 - 62,000 years.


Willard Libby, Nobel Prize in Chemistry (1949)

plant or animal alive : exchanging carbon with its surroundings → same proportion of  $^{14}\text{C}/^{12}\text{C}$  as the biosphere.

Once it dies  $^{14}\text{C}$  it contains decays,  $^{14}\text{C}/^{12}\text{C}$  gradually reduce.


**A mammoth was found in the Siberian permafrost. The  $^{14}\text{C}$  content in the body was only 21 % of that found in living animals. Their  $^{14}\text{C}/^{12}\text{C}$  ratio is  $10^{-12}$ . How old is the mammoth ? The half-life of the radiocarbon is 5730 y.**

# Decay chains



$$A_Y = \lambda_Y N_Y = A_{X,0} \frac{\lambda_Y}{\lambda_Y - \lambda_X} \left( e^{-\lambda_X t} - e^{-\lambda_Y t} \right)$$


relation of  $\lambda_A$  and  $\lambda_B$  ?





$$T_{1/2,X} \gg T_{1/2,Y}$$

$$T_{1/2,X} = 8 \cdot 10^7 \text{h}$$

$$T_{1/2,Y} = 0,8 \text{h}$$







When former Russian spy Alexander Litvinenko died from polonium-210 poisoning several years ago in London, it triggered a murder investigation that developed like a thriller.

Po-210 generate much heat as the atoms decay - it was used in Russian lunar landers to keep the craft's instruments warm at night.

$^{210}\text{Po}$  is an  $\alpha$ -emitter, that has a half-life of 138.4 days,  $E_{\alpha} = 5.3 \text{ MeV}$

# Interaction of the radiation with the matter

## Particles/photons

	I.	II.	III.
a	b		
p	$e^+$	n	$\gamma$
$\alpha$	$e^-$		X

## Partners

1. Electromagnetic field
2. Electron
3. Field of the nucleus
4. Nucleus

## Mechanism

## Effect on

### radiation

### matter

A) Absorption

$\Delta I$

$E_{kin}, E^*$

B) Coherent scattering (only the direction is altered))

$\Delta I$

-

C) Incoherent scattering (also exchange of  $E$ )  
 elastic (no excitation)  
 inelastic


$\Delta I, \Delta E$

$E_{kin}$   
 $E_{kin}, E^*$  27

# 1. Ionizing radiations

# The first step of the ionizing radiation in the matter:

## 1. Neutral excitation



## 2. External ionization




## 3. Internal ionization



## 4. Bremsstrahlung (braking radiation)



# Quantitative description of the interaction



$$v = \sigma n x \rho_A \quad \text{cross section}$$

$$-dn = \sigma(E) n \rho_A dx$$

$$n = n_0 e^{-\sigma(E) \rho_A x}$$

$$I = \frac{n}{t}$$

$$I = I_0 e^{-\mu x} \quad \text{linear absorption coefficient}$$

$$I = I_0 e^{-\mu x} = I_0 e^{-\frac{\mu}{\rho} x \cdot \rho} = I_0 e^{-\mu_m d} \quad \text{mass absorption coefficient}$$

$$x_{1/2} = \frac{\ln 2}{\mu} \quad d_{1/2} = \frac{\ln 2}{\mu_m}$$

# $\alpha$ -radiation

Heavy, charged, high energy

With electrons: incoherent scattering


ionisation and excitation (50-50 %)

$E$  and direction of the alpha particles is modified

With the nucleus: Rutherford-scattering

nuclear reaction (see later)

**! Bremsstrahlung (continuous energy gamma radiation)!**



# $\beta$ -radiation small, charged, limited energy

With electron: incoherent scattering

ionisation (external and internal)


excitation

$E$  and the direction of the radiation changes

$$\frac{\left(\frac{dE}{dx}\right)_r}{\left(\frac{dE}{dx}\right)_{\text{ion}}} = \frac{EZ}{800}$$

With the field of the nucleus: incoherent scattering

! Bremsstrahlung !



$$I = I_0 e^{-\mu'x} = I_0 e^{-\mu d}$$

Linear/mass absorption coefficient



Calculate the activity of 1 kg KCl. 0.012 % of the K atoms is radioactive  $^{40}\text{K}$ . The half life of  $^{40}\text{K}$  is  $1.13 \cdot 10^9$  years.

We prepared a  $^{35}\text{S}$  labelled protein at 12:00, 10 September 2014. The half life of the pure  $\beta^-$  emitter is 88 days. This sample was measured at noon on 26 September and the intensity was found 7000 imp/s. The overall efficiency of the measurement was 22 %. Calculate the activity of the sample in the time of synthesis.


The linear absorption coefficient of gamma radiation of 660 keV in aluminum is  $3,4 \text{ cm}^{-1}$ . Calculate the half thickness. How efficiently will attenuate this radiation an 10 cm aluminum wall ?

# $\gamma$ -radiation

# electromagnetic radiation

## 1. Compton-scattering



Elastic collision of the photon with an electron



$$\mu_C = \frac{\mu'_C}{\rho} = \sigma_C \frac{\rho_A}{\rho} = \sigma_C \frac{N_A Z}{A}$$

$$\sigma_C = \sigma_s + \sigma_a$$


## 2. Photoelectric effect



$$\sigma_f \approx \text{konst.} \frac{Z^2}{(h\nu)^3}$$

$$n(E) = 4 - 5$$



### 3. Pair production



$$\sigma_p = K(E_\gamma - 1,02)^{2,2} Z^2$$

$$I = I_0 e^{-\mu d} = I_0 e^{-(\mu_C + \mu_f + \mu_p)d}$$

## Germanium




## 2. Nuclear reactions

# Cross section (~probability)

## Conventional equation



Transition state



- |    |  |
|----|--|
| 1. | (n, $\gamma$ )   |
|    | (n, f) ${}^{233}\text{U}$ , ${}^{235}\text{U}$ , ${}^{239}\text{Pu}$ , ${}^{241}\text{Pu}$ |
|    | ${}^{10}\text{B}(n, \alpha)$   |
|    | ${}^6\text{Li}(n, \alpha)$   |
| 2. | ( $\gamma$ , n)  |
|    | (n, 2n)  |
|    | (n, $\alpha$ )   |
|    | (p, )  |
|    | (d, )  |

Tunnel effect


# Kinetics of the nuclear reactions

$$\frac{dN^*}{dt} = \sigma_a N \phi - \lambda N^*$$

$$N^* = N_{\infty}^* [1 - \exp(-\lambda t)]$$


$$A = A_{\infty} [1 - \exp(-\lambda t)]$$

$$A_{\infty} = \lambda N_{\infty}^* = \phi \sigma_a N$$



$$A' = \lambda N^* =$$

$$= A_{\infty} [1 - \exp(-\lambda t)] \exp(-\lambda t_h)$$





We intend to obtain  $^{65}\text{Ni}$  with neutron irradiation. Therefore, we expose 1 g of Ni (with a  $^{64}\text{Ni}$  content of 91 %) to neutrons with a flux  $\Phi=10^{12}$  1/cm<sup>2</sup>s. The cross section  $\sigma$  of the



reaction is  $1.55 \cdot 10^{-28}$  m<sup>2</sup>. The half-life of  $^{65}\text{Ni}$  is 2.52 h.

- i) How long should the irradiation last if we want to reach 80 % of the saturation activity?
- ii) Estimate the ratio of the  $^{64}\text{Ni}/^{65}\text{Ni}$  isotopes in the sample after being „cooled“ for the same period as the activation lasted.

# Interaction of neutrons with the matter

relatively heavy, no charge, energy ?

- elastic scattering

Table R8. The energy absorption efficiency of light elements

$$(E_0 = 2 \text{ MeV}, E = kT)$$

Element	$\Delta\bar{E}$ , keV	$n$
$^1\text{H}$	1000	18
$^2\text{D}$	888	24
$^4\text{He}$	640	41
Be	360	50
C	284	111
Al	137	240

- inelastic scattering

Excited nucleus,  $h\nu$

- neutron capture

(absorption): (n,?)

## Due to the strong $E$ dependence,

### 1. Slow

a) cold

$$E < 0.025 \text{ eV}$$

b) *thermal*

$$0.025 \text{ eV} < E < 0.44 \text{ eV}$$

c) resonance

$$0.44 \text{ eV} < E < 1000 \text{ eV}$$

### 2. Medium

$$1 \text{ keV} < E < 500 \text{ keV}$$

### 3. Fast

$$0.5 \text{ MeV} < E < 10 \text{ MeV}$$


### 4. High energy

$$10 \text{ MeV} < E < 50 \text{ MeV}$$

### 5. Super fast

$$50 \text{ MeV} < E$$




# Examples of practical relevance




## $(n, f)$ fission

Fuel	Source of the fuel	Neutron energy needed
$^{235}\text{U}$	natural uranium	$0.025 \text{ eV} < E_{\text{neutron}} < 0.44 \text{ eV}$ (thermal)
$^{233}\text{U}$	from thorium with n absorption	$0.025 \text{ eV} < E_{\text{neutron}} < 0.44 \text{ eV}$ (thermal)
$^{239}\text{Pu}$	from $^{238}\text{U}$ with n absorption	$0.025 \text{ eV} < E_{\text{neutron}} < 0.44 \text{ eV}$ (thermal)
$^{241}\text{Pu}$	from $^{238}\text{U}$ with n absorption	$0.025 \text{ eV} < E_{\text{neutron}} < 0.44 \text{ eV}$ (thermal)
$^{238}\text{U}$	natural uranium	$0.5 \text{ MeV} < E_{\text{neutron}} < 10 \text{ MeV}$ (fast)
$^{232}\text{Pu}$	natural thorium	$0.5 \text{ MeV} < E_{\text{neutron}} < 10 \text{ MeV}$ (fast)

# Fission (n, f)





50 ways, 300 isotopes 35 elements



# Distribution 200 MeV

- kinetic energy of fission products:  $\approx 160 \text{ MeV}$
- kinetic energy of the neutrons:  $\approx 5 \text{ MeV}$
- energy of the  $\gamma$ -rays:  $\approx 5 \text{ MeV}$
- energy of the secondary radioactive decay:  $\approx 20 \text{ MeV}$
- energy released at neutron capture:  $\approx 10 \text{ MeV}$




**Self-sustaining chain reaction: control**

# Detection of nuclear radiations




# Interaction with matter: Linear energy transfer (LET)

air



Path



$$dE / dx \approx 1/v^2$$

# The first step of the ionizing radiation in the matter:

## 1. Neutral excitation



## 2. External ionization



## 3. Internal ionization



## 4. Bremsstrahlung (braking radiation)



What do we want to know?

yes/no

type of radiation

energy of radiation

source

activity  $(I=k\eta A)$


integral

real time evaluation


delayed evaluation

rate


# Geiger-Müller (GM) counter (gas ionisation detector)



Dead time



Characteristic curve



# Semiconductor detectors

Typical semiconductors


	Si	Ge	CdTe
Atomic number, $Z$	14	32	48 - 52
Energy gap, eV	1.12	0.74	1.47
Ionisation energy, eV	3.61	2.98	4.43

Ge(Li)

HPGe, Si(Li)

# Scintillation detectors

Scintillator (material depends on the radiation) + photomultiplier



# Typical scintillation crystals

Depends on the type of radiation

NaI(Tl)                      gamma

Plastic                      beta


ZnS                      alpha

Liquid scintillation technique

for low  $E$  isotopes ( $^3\text{H}$ ,  $^{14}\text{C}$ )

scintillator and radioactive material dissolved  
in the same solution

# Comparison of a scintillation and a semiconductor spectrum





## Comparison of the features of the main detector types

<b>Properties</b>	<b>GM counter</b>	<b>Scintillation detector</b>	<b>Semiconductor detector</b>
<b>Field of application</b>	Primarily for particle radiation measurements	Measurements of any radioactive radiation types	Measurements of any radioactive radiation
<b>Measurement efficiency</b>	For particle radiation ( $\alpha$ , $\beta$ , $n$ ) near 100% for electromagnetic radiation 1 or 2%	Generally good	Generally good strongly temperature dependent at some types
<b>Dead time</b>	< 1 ms	<1 $\mu$ s	<0.1 $\mu$ s
<b>Energy selectivity (qualitative identification of the radioactive source)</b>	Non-selective	Selective	Very selective
<b>Costs</b>	Low	High, due to accessories	High
<b>Other aspects</b>	Limited but usually long life time	High counting rates	For drifted semiconductors, cooling required both for measurement and storage